

This is a repository copy of *Recurrence to shrinking targets on typical self-affine fractals*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/112732/>

Version: Accepted Version

---

**Article:**

Koivusalo, Henna and Ramírez, Felipe A. (2018) Recurrence to shrinking targets on typical self-affine fractals. *Proceedings of the Edinburgh Mathematical Society*. pp. 387-400. ISSN 1464-3839

<https://doi.org/10.1017/S0013091517000256>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# RECURRENCE TO SHRINKING TARGETS ON TYPICAL SELF-AFFINE FRACTALS

HENNA KOIVUSALO AND FELIPE A. RAMÍREZ

**ABSTRACT.** We explore the problem of finding the Hausdorff dimension of the set of points that recur to shrinking targets on a self-affine fractal. To be exact, we study the dimension of a certain related symbolic recurrence set. In many cases this set is equivalent to the recurring set on the fractal.

## 1. INTRODUCTION

Given a dynamical system  $(X, T)$ , and a sequence  $\mathcal{B} = (B_k)$  of subsets of  $X$ , a **shrinking target problem** is the study of the size of the recurring set

$$R(\mathcal{B}) = \{x \in X : T^k(x) \in B_k \text{ for infinitely many } k \in \mathbb{N}\}.$$

Viewing the sets  $B_k$  as “targets,” the set  $R(\mathcal{B})$  is the set of points whose orbits under the dynamics hit the targets infinitely often.

Interest in recurrence is classical in the study of dynamical systems. There are many ways to interpret the “size” of  $R(\mathcal{B})$ . One way is with respect to an invariant measure on  $X$ . In this context, for example, for any positive measure set  $B \subset X$ , the Poincaré recurrence theorem implies that almost all points of  $X$  return to  $B$  infinitely often, so that  $R(\mathcal{B})$  has full measure when  $B_k = B$  for all  $k$ . It is thus fairly natural to ask what happens when the targets shrink. In many settings  $R(\mathcal{B})$  has either full or zero measure depending on the rate of shrinking of  $B_k$ . See, for example, Chernov and Kleinbock [5] for a general treatment of such a problem.

For exponentially decreasing balls on Julia sets, Hill and Velani calculated first the Hausdorff dimension [10] and then the Hausdorff measure [12] of the recurring set, in the latter case proving a 0-1-law for  $R(\mathcal{B})$ . Further, in [11] they considered the dimension of  $R(\mathcal{B})$  for toral automorphisms. More recently, the problem has been studied for infinitely branching Markov maps [24], piecewise expanding maps of the

---

*Date:* October 20, 2015.

H.K. was supported by EPSRC Programme Grant EP/L001462 and F.A.R. by EP/J018260/1.

unit interval [22] and in the dynamical system of continued fraction expansions [20]. In all of these cases the dimension of the recurring set is given as a zero point of a generalized pressure functional.

Recurring points are not only an interesting object in abstract settings; they also appear in number theory in the study of Diophantine approximation. The doubly-metric inhomogeneous version of Khintchine's Theorem, usually stated as a 0-1-law for the set of "approximable" pairs in  $\mathbb{R}^d \times \mathbb{R}^d$ , can be interpreted through Fubini's Theorem as a 0-1-law for the recurring set of shrinking targets on a torus, which holds for almost any toral translation. This connection to toral translations is made pleasantly visible by Fayad [8], where among other things he gives a proof of a result of Kurzweil [19] from this point-of-view. Similarly, one can interpret the classical Jarník-Besicovitch theorem as a dimension result for a recurring set of shrinking targets on the torus. For a more recent example, see Bugeaud, Harrap, Kristensen and Velani [4].

The above results mainly focus on conformal dynamical systems. In this note we study the shrinking target problem in a simple non-conformal case, namely, under affine dynamics. In particular, we will calculate the dimension of the recurring set on a certain fractal set, the definition of which will be given next.

Hutchinson [13] originated the systematic study of iterated function systems (IFSs)  $\{f_1, \dots, f_m\}$ , which define a fractal set through the identity  $E = \cup_{i=1}^m f_i(E)$ . In particular, the **self-affine fractal** is the unique, nonempty, compact set  $E$  satisfying

$$E_{(a_1, \dots, a_m)} = E = \bigcup_{i=1}^m T_i(E) + a_i = \bigcup_{i=1}^m f_i(E),$$

generated by iterating the affine maps  $\{f_i = T_i + a_i\}_{i=1, \dots, m}$ . Here  $\{T_1, \dots, T_m\}$  is a collection of bijective linear contractions on  $\mathbb{R}^d$  and  $a_1, \dots, a_m \in \mathbb{R}^d$  are translation vectors. If the images  $f_i(E)$  are disjoint it is possible to define a mapping  $F : E \rightarrow E$  by setting

$$F(x) = f_i^{-1}(x)$$

for points  $x \in f_i(E)$ , for all  $i = 1, \dots, m$ . This sets up a dynamical system  $(E, F)$ , and so it makes sense to formulate a shrinking target problem on  $E$ .

The study of dimensional properties of self-affine sets comes in three branches: First, as self-affine carpets due to, originally, Bedford [1] and McMullen [21]; second, for generic self-affine sets in the footsteps of Falconer [7]; third, for *every* self-affine set in some family,

as in [6]. In this paper we consider a shrinking target problem associated to the set  $E$  in the generic sense, that is, for generic translations  $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{R}^{dm}$ . A complementary problem of mass escape for carpet type sets has been studied by Ferguson, Jordan and Rams in [9].

The main result of this article can be stated informally as follows.

**Main Theorem** (An informal version of Theorem 1). *The dimension of the recurring set to shrinking targets on self-affine fractals is almost surely the unique zero point  $s_0 \in \mathbb{R}^+$  of a modified pressure function  $P(s)$ .*

We will impose a quasimultiplicativity condition (6) on the maps  $T_i$ , and give the modified pressure formula (3). In this article, the target sets  $B_k$  are dynamical balls corresponding to cylinder sets in the symbolic space (for definitions, see §2.1). These target sets match the dynamics on the set. The method of proof is largely classical and the main difficulty lies in finding the right definition for the pressure functional. Also, in order to obtain the lower bound we need to construct a Cantor type subset of the recurring set. We do this in the spirit of the proof of the mass transference principle of Beresnevich and Velani [2]. Their result cannot be applied to the affine setting directly.

It is classical to study the size of the recurring set in terms of invariant measures, and it turns out that, at least in certain cases, this problem is easily solved based on [5]. In particular, in §6 we notice that as a corollary of [5, Theorem 2.1] it is possible to measure the size of the recurring set in terms of the Gibbs measure at the dimension of the self-affine set  $E$ . According to [15] Gibbs measures are measures of maximal dimension on  $E$ , which in part motivates their use in this context. For further reading on general thermodynamic formalism and Gibbs measures we refer to [3], and in relation to self-affine sets, to [15]).

**Corollary to [5]** (An informal version of Corollary 8). *In terms of the Gibbs measure at the dimension of the self-affine set  $E$ , either almost all or almost no points are recurring, depending on whether the sum of the measures of the target sets diverges or converges.*

Even though these formulations of the shrinking target problem give a very natural starting point to the analysis of shrinking targets for nonconformal dynamics, some questions remain. We discuss these at the end of §6.

**Acknowledgments.** We are grateful to Thomas Jordan for suggesting the problem to the first author, and for helpful discussions. We also

thank Antti Käenmäki for taking the time to read our manuscript, and for giving valuable feedback.

## 2. SHRINKING TARGETS, FRACTALS, AND SYMBOLIC DYNAMICS

Let  $E_a$ ,  $f_i$ ,  $T_i$  be as in the introduction.

**2.1. Self-affine fractals and symbolic dynamics.** It is standard to associate to  $E_a$  a symbolic dynamical system given by the left shift  $\sigma$  on the sequence space  $\mathcal{A}^{\mathbb{N}} = \{1, \dots, m\}^{\mathbb{N}}$ . One identifies points  $\mathbf{i} = (i_1, i_2, \dots) \in \mathcal{A}^{\mathbb{N}}$  with points on  $E_a$  through the projection

$$\pi(\mathbf{i}) = \lim_{k \rightarrow \infty} f_{i_1} \circ \dots \circ f_{i_k}(0) \in E_a,$$

and if this projection is injective, then  $\sigma$  induces a well-defined dynamical system on  $E_a$ , taking the point  $\pi(\mathbf{i}) = p \in E_a$  to  $f_{i_1}^{-1}(p)$ . In this case, the shrinking targets problem (defined in the next section) on the symbolic dynamical system corresponds to a shrinking targets problem on the fractal with the dynamics induced by  $\sigma$ .

**2.2. Shrinking targets on a symbolic dynamical system.** A natural way to define a shrinking targets problem on the dynamical system  $\sigma : \mathcal{A}^{\mathbb{N}} \rightarrow \mathcal{A}^{\mathbb{N}}$  is to select a target point  $\mathbf{j} \in \mathcal{A}^{\mathbb{N}}$  and define the targets to be cylinder sets

$$[\mathbf{j}|_k] = \{\mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \mathbf{i}|_k = \mathbf{j}|_k\} \subset \mathcal{A}^{\mathbb{N}}$$

of increasing length  $k \in \mathbb{N}$  (and hence decreasing diameter) around that point, where  $\mathbf{j}|_k$  and  $\mathbf{i}|_k$  denotes the truncations  $(j_1, \dots, j_k)$  and  $(i_1, \dots, i_k)$  of  $\mathbf{j}$  and  $\mathbf{i}$ , respectively.

In this paper we treat the shrinking targets problem for  $\mathcal{B} = \{B_k\}_{k \in \mathbb{N}}$  where  $B_k = [\mathbf{j}|_{\ell_k}]$  and  $\{\ell_k\} \subseteq \mathbb{N}$  is some fixed (non-decreasing) sequence. We denote the recurring set  $R(\mathcal{B})$  by  $R(\mathbf{j})$ . It is exactly

$$\begin{aligned} R(\mathbf{j}) &= \{\mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \sigma^k(\mathbf{i})|_{\ell_k} = \mathbf{j}|_{\ell_k} \text{ for infinitely many } k \in \mathbb{N}\} \\ &= \limsup_{k \rightarrow \infty} R(\mathbf{j}, k) \end{aligned}$$

where  $R(\mathbf{j}, k) = \{\mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \sigma^k(\mathbf{i})|_{\ell_k} = \mathbf{j}|_{\ell_k}\}$ .

**2.3. Shrinking targets on self-affine fractals.** We project our symbolic shrinking targets problem to the fractal  $E_a$  through  $\pi$ . That is, we seek the Hausdorff dimension of the set

$$\tilde{R}(\pi(\mathbf{j})) := \pi(R(\mathbf{j})) \subset E_a.$$

An intuitive interpretation is to think of the shrinking targets around  $\pi(\mathbf{j})$  as the succession of “construction sets” of  $E_a$  to which  $\pi(\mathbf{j})$  belongs. (One may think of the standard Cantor set, for example, and

take the shrinking targets to be subintervals appearing in its standard construction.) Though these are very natural shrinking targets on a fractal, the notion is really only meaningful if the affine maps  $\{f_i\}$  satisfy a strong enough separation condition. Still, the set  $\tilde{R}(\pi(\mathbf{j}))$  is well-defined and can be studied regardless.

### 3. LIST OF NOTATIONS

This list of standard notations may be used as reference.

- $\mathcal{A}^{\mathbb{N}} = \{1, \dots, m\}^{\mathbb{N}}$  is the sequence space on  $m$  letters.
- $\mathbf{j}|_k = j_1 j_2 \dots j_k$  is the string obtained by truncating  $\mathbf{j} \in \mathcal{A}^{\mathbb{N}}$ .
- $|\mathbf{i}|$  is the length of a finite word  $\mathbf{i} = (i_1, \dots, i_k)$ , so for example  $|\mathbf{j}|_k = k$ .
- $\mathbf{i}\mathbf{j}|_k = i_1 i_2 \dots i_{|\mathbf{i}|} j_1 j_2 \dots j_k$  is a concatenation of finite words.
- $\mathbf{i} \wedge \mathbf{i}'$  is the maximal common beginning of  $\mathbf{i}, \mathbf{i}' \in \mathcal{A}^{\mathbb{N}}$ .
- $[\mathbf{q}] = \{\mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \mathbf{i}|_{|\mathbf{q}|} = \mathbf{q}\}$  is the cylinder set defined by the finite word  $\mathbf{q}$ .
- Let  $f_i : \mathbb{R}^d \rightarrow \mathbb{R}^d$  be the affine maps  $f_i(x) = T_i(x) + a_i$ .
- For  $|\mathbf{i}| = k$ , denote  $f_{\mathbf{i}} = f_{i_1} \circ \dots \circ f_{i_k}$ , and similarly for  $T_{\mathbf{i}}$ .
- $R(\mathbf{j}) = \{\mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \sigma^k(\mathbf{i})|_{\ell_k} = \mathbf{j}|_{\ell_k} \text{ infinitely often}\}$  is our symbolic recurring set.
- $\tilde{R}(\pi(\mathbf{j})) = \pi(R(\mathbf{j}))$ .
- $R(k) = R(\mathbf{j}, k) = \{\mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \sigma^k(\mathbf{i})|_{\ell_k} = \mathbf{j}|_{\ell_k}\}$ , so that we have  $R(\mathbf{j}) = \limsup R(k)$ .

### 4. THE SINGULAR VALUE FUNCTION

For a linear map  $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$  define the singular values

$$\sigma_1(T) \geq \dots \geq \sigma_d(T)$$

as the lengths of the semiaxes of the ellipsoid  $T(B(0, 1))$ . For any  $0 \leq t \leq d$ , let

$$\varphi^t(T) = \sigma_1(T) \dots \sigma_{[t]}(T) \sigma_{[t]+1}(T)^{\{t\}}$$

where  $[t]$  is the integer part of  $t$  and  $\{t\}$  is the fractional part. The function  $\varphi^t(T)$  is called the singular value function, and it satisfies

$$(1) \quad \sigma_d(T)^t \leq \varphi^t(T) \leq \sigma_1(T)^t.$$

Furthermore, if  $T_1, \dots, T_n$  are linear maps on  $\mathbb{R}^d$ , then

$$(2) \quad \min_{i=1, \dots, n} \sigma_d(T_i)^\delta \leq \frac{\sum_{i=1}^n \varphi^{t+\delta}(T_i)}{\sum_{i=1}^n \varphi^t(T_i)} \leq \max_{i=1, \dots, n} \sigma_1(T_i)^\delta$$

for all sufficiently small  $\delta > 0$ . The singular value function is submultiplicative, that is, if also  $U : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is a linear map, then  $\varphi^t(TU) \leq \varphi^t(T)\varphi^t(U)$ . For proofs see [7], for example.

We study collections of linear maps for which there exists some constant  $D \in (0, 1)$  such that

$$\varphi^t(T_{\mathbf{i}\mathbf{i}'}) \geq D\varphi^t(T_{\mathbf{i}})\varphi^t(T_{\mathbf{i}'})$$

for any finite words  $\mathbf{i}, \mathbf{i}'$ . This condition holds, for example, for similarities and for maps on  $\mathbb{R}^2$  satisfying the *cone condition*, meaning that there is a cone that is mapped strictly into itself by all the  $T_i$ 's, see Käenmäki and Shmerkin [16].

## 5. THE MAIN RESULT AND ITS PROOF

Given the affine maps  $\{f_i\}_{i=1,\dots,m}$ , a point  $\mathbf{j} \in \mathcal{A}^{\mathbb{N}}$ , and  $s \in \mathbb{R}^+$ , we define

$$(3) \quad P(s, \mathbf{j}) := \lim_{k \rightarrow \infty} \frac{1}{k} \log \sum_{|\mathbf{i}|=k} \varphi^s(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}).$$

This is a modification of the standard pressure function (see [7] for example), which is defined to be

$$(4) \quad \mathcal{P}(s) = \lim_{k \rightarrow \infty} \frac{1}{k} \log \sum_{|\mathbf{i}|=k} \varphi^s(T_{\mathbf{i}}).$$

In fact, notice<sup>1</sup> that in the presence of (6) we have

$$(5) \quad P(s, \mathbf{j}) = \mathcal{P}(s) + \lim_{k \rightarrow \infty} \frac{1}{k} \log \varphi^s(T_{\mathbf{j}|_{\ell_k}}).$$

Notice also that it is possible for this limit to not exist. However, if it *does* exist for some  $\mathbf{j}$  and  $s \in \mathbb{R}^+$ , then in fact  $P(s, \mathbf{j})$  is a continuous and non-increasing function of  $s$ . In this case, we prove that the Hausdorff dimension of the recurring set  $\tilde{R}(\pi(\mathbf{j}))$  is almost surely the unique zero point  $s_0 \in \mathbb{R}^+$  of  $P(\cdot, \mathbf{j})$ .

**Theorem 1.** *Let  $E_{\mathbf{a}}$  be the fractal determined by the affine maps  $\{T_i + a_i\}_{i=1}^m$ , and suppose there is some  $D \in (0, 1)$  such that*

$$(6) \quad \varphi^t(T_{\mathbf{i}\mathbf{i}'}) \geq D\varphi^t(T_{\mathbf{i}})\varphi^t(T_{\mathbf{i}'})$$

*for any finite words  $\mathbf{i}, \mathbf{i}'$ . Furthermore, assume that  $\max_i \sigma_1(T_i) < \frac{1}{2}$ . Consider shrinking targets defined by cylinder sets of length  $\{\ell_k\}_{k \in \mathbb{N}} \subseteq \mathbb{N}$  around some fixed  $\mathbf{j} \in \mathcal{A}^{\mathbb{N}}$ .*

---

<sup>1</sup>We thank an anonymous referee for making this observation.

If  $\lim_{k \rightarrow \infty} \frac{1}{k} \log \varphi^s(T_{\mathbf{j}|_{\ell_k}})$  exists in  $[-\infty, 0]$ , then with respect to any ergodic  $\sigma$ -invariant probability measure there is almost surely a unique number  $s_0 \in \mathbb{R}^+$  satisfying  $P(s_0, \mathbf{j}) = 0$ , and for almost every  $\mathbf{a} \in \mathbb{R}^{dm}$ , the Hausdorff dimension of the recurring set  $\tilde{R}(\pi(\mathbf{j})) \subset E_{\mathbf{a}}$  equals  $s_0$ .

*Remark.* If the maps  $\{T_i + a_i\}_{i=1}^m$  satisfy the strong separation condition then the set  $\tilde{R}(\mathbf{a}, \mathbf{j})$  from Theorem 1 has a dynamical meaning as a subset of the fractal  $E_{\mathbf{a}}$ .

Of course, it now becomes a question of *when*  $\lim_{k \rightarrow \infty} \frac{1}{k} \log \varphi^s(T_{\mathbf{j}|_{\ell_k}})$  exists. In the following corollary we show that in particular it exists if  $\lim_{k \rightarrow \infty} \ell_k/k$  does.

**Corollary 2.** *With the assumptions and notation of Theorem 1, if  $\lim_{k \rightarrow \infty} \ell_k/k$  exists, then Theorem 1 applies and we will have*

$$(7) \quad \dim_{\mathcal{H}} \tilde{R}(\pi(\mathbf{j})) = \begin{cases} \dim_{\mathcal{H}} E_{\mathbf{a}} & \text{if } \lim_{k \rightarrow \infty} \ell_k/k = 0 \\ s_0 & \text{if } \lim_{k \rightarrow \infty} \ell_k/k \in (0, \infty) \\ 0 & \text{if } \lim_{k \rightarrow \infty} \ell_k/k = \infty \end{cases}$$

*almost surely (in the sense of the theorem).*

*Proof.* If  $\lim_{k \rightarrow \infty} \ell_k/k = \infty$ , then the result follows from Lemma 5, and in fact holds for *every* target point  $\mathbf{j}$  and translation  $\mathbf{a}$ .

If  $\lim_{k \rightarrow \infty} \ell_k/k \in [0, \infty)$ , then Lemma 3 implies the existence of  $\lim_{k \rightarrow \infty} \frac{1}{k} \log \varphi^s(T_{\mathbf{j}|_{\ell_k}})$ , so Theorem 1 applies. It is only left to observe that in the case where  $\ell_k = o(k)$ , we will have  $\mathcal{P}(s) = P(s, \mathbf{j})$ , so the unique zero points coincide, and therefore the recurring set has the same Hausdorff dimension as the fractal.  $\square$

**5.1. Existence and uniqueness.** The next lemma shows that if there is some  $L \in [0, \infty)$  for which  $\ell_k \sim L \cdot k$ , then there exists a unique number  $s_0 \in \mathbb{R}^+$  such that  $P(s_0, \mathbf{j}) = 0$ . It is furthermore almost surely independent of  $\mathbf{j}$  with respect to any  $\sigma$ -invariant measure.

**Lemma 3.** *With the same assumptions and notation as in Theorem 1, suppose that  $\lim_{k \rightarrow \infty} \ell_k/k \in [0, \infty)$ . Then, with respect to any  $\sigma$ -ergodic probability measure on  $\mathcal{A}^{\mathbb{N}}$ , there almost surely exists a number  $s_0 \in \mathbb{R}^+$  satisfying  $P(s_0, \mathbf{j}) = 0$ . Furthermore, this number is unique and almost surely independent of  $\mathbf{j} \in \mathcal{A}^{\mathbb{N}}$ .*

*Proof.* Suppose  $\ell_k \sim L \cdot k$ ,  $L > 0$ . Fix  $s$ . Let

$$X(m, n) = \log \varphi^s(T_{\sigma^m(\mathbf{j})|_{n-m}}).$$

We will apply Kingman's subadditive ergodic theorem to the sequence  $X(m, n)$  with respect to the transformation  $\sigma$ , so we must check that



$X$  is subadditive and  $\sigma$ -equivariant. Clearly all  $X(m, n)$  are integrable, since they only depend on finite words.

**Equivariance** is verified by the following routine calculation:

$$\begin{aligned} X(m, n) \circ \sigma &= \log \varphi^s(T_{\sigma^m(\sigma(\mathbf{j}))|_{n-m}}) \\ &= \log \varphi^s(T_{\sigma^{(m+1)}(\mathbf{j})|_{(n+1)-(m+1)}}) = X(m+1, n+1). \end{aligned}$$

For **subadditivity** we must check that  $X(0, n) \leq X(0, m) + X(m, n)$ . Again, this is routine:

$$\begin{aligned} X(0, m) + X(m, n) &= \log \varphi^s(T_{\mathbf{j}|_m}) + \log \varphi^s(T_{\sigma^m(\mathbf{j})|_{n-m}}) \\ &= \log \varphi^s(T_{\mathbf{j}|_m}) \varphi^s(T_{\sigma^m(\mathbf{j})|_{n-m}}) \geq \log \varphi^s(T_{\mathbf{j}|_n}) = X(0, n). \end{aligned}$$

Therefore, by the subadditive ergodic theorem, there almost surely exists a limit

$$\lim_{n \rightarrow \infty} \frac{X(0, n)}{n} = \lim_{k \rightarrow \infty} \frac{X(0, \ell_k)}{\ell_k}$$

which is  $\sigma$ -invariant and so almost everywhere constant in  $\mathbf{j}$  with respect to any  $\sigma$ -ergodic measure. Hence,

$$\lim_{k \rightarrow \infty} \frac{X(0, \ell_k)}{k} = \lim_{k \rightarrow \infty} \frac{X(0, \ell_k)}{\ell_k} \cdot \frac{\ell_k}{k} = L \lim_{k \rightarrow \infty} \frac{X(0, \ell_k)}{\ell_k}$$

also exists and is almost surely constant in  $\mathbf{j}$ , and similarly for

$$P(s, \mathbf{j}) = \mathcal{P}(s) + \lim_{k \rightarrow \infty} \frac{X(0, \ell_k)}{k}.$$

Notice that the limit is continuous and strictly decreasing in  $s$  since by (2),

$$\sigma_-^{(k+\ell_k)\delta} \leq \frac{\sum_{|\mathbf{i}|=k} \varphi^{s+\delta}(T_{\mathbf{i}\mathbf{j}|_{\ell_k}})}{\sum_{|\mathbf{i}|=k} \varphi^s(T_{\mathbf{i}\mathbf{j}|_{\ell_k}})} \leq \sigma_+^{(k+\ell_k)\delta},$$

where  $0 < \sigma_- = \min_i \sigma_d(T_i)$  and  $\sigma_+ = \max_i \sigma_1(T_i) < 1$ .

Since  $P(0, \mathbf{j}) > 0$  and  $P(s, \mathbf{j}) \rightarrow -\infty$  when  $s \rightarrow \infty$ , the existence of a unique zero follows.  $\square$

**5.2. Upper bound.** The following lemma shows that  $s_0$  is an upper bound for the dimension of the recurring set  $\tilde{R}(\pi(\mathbf{j}))$ .

**Lemma 4.** *Let  $\{\ell_k\} \subseteq \mathbb{N}$  be a non-decreasing divergent sequence and suppose that  $P(\cdot, \mathbf{j})$  is decreasing, with unique zero  $s_0 \in \mathbb{R}$ . Then the Hausdorff dimension of  $\tilde{R}(\pi(\mathbf{j}))$  is bounded above by  $s_0$ .*

*Proof.* We will show that for any  $t > s_0$ , we have  $\mathcal{H}^t(\tilde{R}(\pi(\mathbf{j}))) = 0$ , where  $\mathcal{H}^t$  denotes Hausdorff measure.

Notice that when  $t > s_0$ , then for all large  $k$ ,

$$\sum_{|\mathbf{i}|=k} \varphi^t(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}) \leq \exp(-k\varepsilon)$$

for some  $\varepsilon > 0$ . Therefore

$$(8) \quad \sum_{k=0}^{\infty} \sum_{|\mathbf{i}|=k} \varphi^t(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}) < \infty.$$

For all  $N$ , we have that

$$R(\mathbf{j}) = \limsup_{k \rightarrow \infty} \{ \mathbf{i} \in \mathcal{A}^{\mathbb{N}} \mid \sigma^k(\mathbf{i})|_{\ell_k} = \mathbf{j}|_{\ell_k} \} \subset \bigcup_{k=N}^{\infty} \bigcup_{|\mathbf{i}|=k} [\mathbf{i}\mathbf{j}|_{\ell_k}],$$

hence,

$$\tilde{R}(\pi(\mathbf{j})) \subset \bigcup_{k=N}^{\infty} \bigcup_{|\mathbf{i}|=k} \pi[\mathbf{i}\mathbf{j}|_{\ell_k}].$$

Since the  $T_i$ 's are contractions, there is some  $M > 0$  sufficiently large that  $f_i(B(0, M)) \subset B(0, M)$  for all  $i = 1, \dots, m$ . Then we have the inclusion  $\pi[\mathbf{i}\mathbf{j}|_{\ell_k}] \subset f_{\mathbf{i}\mathbf{j}|_{\ell_k}}(B(0, M))$ , and therefore

$$\tilde{R}(\pi(\mathbf{j})) \subset \bigcup_{k=N}^{\infty} \bigcup_{|\mathbf{i}|=k} f_{\mathbf{i}\mathbf{j}|_{\ell_k}}(B(0, M)).$$

Recall that  $\sigma_1(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}), \dots, \sigma_d(T_{\mathbf{i}\mathbf{j}|_{\ell_k}})$  are the lengths of the semiaxes of the ellipsoid  $T_{\mathbf{i}\mathbf{j}|_{\ell_k}}(B(0, 1))$ , in decreasing order. Therefore, the number of cubes of side-length  $\sigma_{\lfloor t \rfloor + 1} q(T_{\mathbf{i}\mathbf{j}|_{\ell_k}})$  required to cover the sets  $f_{\mathbf{i}\mathbf{j}|_{\ell_k}}(B(0, M))$  is bounded by

$$A \sigma_1(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}) \dots \sigma_{\lfloor t \rfloor + 1}(T_{\mathbf{i}\mathbf{j}|_{\ell_k}})^{-\lfloor t \rfloor},$$

where  $A > 0$  is some constant. We can bound the Hausdorff measure

$$\begin{aligned} \mathcal{H}^t(\tilde{R}(\pi(\mathbf{j}))) &\leq A \sum_{k=N}^{\infty} \sum_{|\mathbf{i}|=k} \sigma_1(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}) \dots \sigma_{\lfloor t \rfloor + 1}(T_{\mathbf{i}\mathbf{j}|_{\ell_k}})^{-\lfloor t \rfloor} \cdot \sigma_{\lfloor t \rfloor + 1}(T_{\mathbf{i}\mathbf{j}|_{\ell_k}})^t \\ &= A \sum_{k=N}^{\infty} \sum_{|\mathbf{i}|=k} \varphi^t(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}). \end{aligned}$$

But  $N \in \mathbb{N}$  was arbitrary, so recalling (8), we can take  $N \rightarrow \infty$  to see that  $\mathcal{H}^t(\tilde{R}(\pi(\mathbf{j}))) = 0$ . This proves the lemma.  $\square$

The next lemma shows that if the shrinking targets are projected cylinder sets whose lengths grow super-linearly, then the corresponding recurring set is zero-dimensional.

**Lemma 5.** *Suppose that  $\lim_{k \rightarrow \infty} \ell_k/k = \infty$ . Then  $\dim_{\mathcal{H}} \tilde{R}(\pi(\mathbf{j}), \ell_k) = 0$ .*

*Proof.* We show that  $\mathcal{H}^\delta(\tilde{R}(\pi(\mathbf{j}), \ell_k)) = 0$  for any  $\delta > 0$ .

Let  $g(k) = \ell_k/k$ . Then  $g(k) \rightarrow \infty$  as  $k \rightarrow \infty$ , and

$$\sum_{k \in \mathbb{N}} \sum_{|\mathbf{i}|=k} \varphi^\delta(T_{\mathbf{i}\mathbf{j}}|_{\ell_k}) \leq \sum_{k \in \mathbb{N}} m^k (\sigma_+^\delta)^{k+\ell_k} \leq \sum_{k \in \mathbb{N}} \left( m \cdot \sigma_+^{\delta(1+g(k))} \right)^k < \infty$$

because for all  $k$  large enough, the parenthetical quantity is  $< 1/2$  (say). The argument from the proof of Lemma 4 shows that  $\mathcal{H}^\delta(\tilde{R}(\pi(\mathbf{j}))) = 0$ , and proves the lemma.  $\square$

**5.3. Lower bound.** Fix numbers  $s < t < s_0$ . Let  $(n_k)$  be a sequence of natural numbers. Let  $C = \bigcap_{k=1}^{\infty} R(n_k)$ , where

$$\begin{aligned} R(n_k) &= \{ \mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \sigma^{n_k}(\mathbf{i})|_{\ell_{n_k}} = \mathbf{j}|_{\ell_{n_k}} \} \\ &= \{ \mathbf{i} \in \mathcal{A}^{\mathbb{N}} : \sigma^{n_k}(\mathbf{i}) \in B_{n_k} \}. \end{aligned}$$

Notice that  $C(k) = R(n_k)$  are finite unions of compact cylinder sets and thus compact, so that  $C$  is compact as well. Furthermore,  $C \subset R = \limsup R(k)$ , and we hope to prove that  $\dim_{\mathcal{H}} C \geq s$ .

**Lemma 6.** *Let  $E_a$  and  $\{\ell_k\}$  be as in Theorem 1. Fix  $t < s_0$ . Then there is a probability measure  $\mu$  supported on  $C$ , and a constant  $H$ , satisfying*

$$(9) \quad \mu[\mathbf{q}] \leq H \varphi^t(T_{\mathbf{q}})$$

for all finite words  $\mathbf{q}$ .

*Remark.* A measure  $\mu$  satisfying (9) could possibly be found through a construction similar to one used in [23], or perhaps an adaptation of [14], and this may relieve us from having to assume quasi-multiplicativity (6). However, quasi-multiplicativity is also needed in the observation (5). For this reason, we have opted to present a self-contained proof of the existence of  $\mu$ , even if our proof *does* rely on (6).

*Proof.* Let  $(n_k)$  be a sequence of natural numbers to be fixed later, with the property that  $n_k \geq n_{k-1} + \ell_{n_{k-1}}$  for all  $k \in \mathbb{N}$ . Let

$$\mu_k = \frac{\sum_{\mathbf{i} \in A(k)} \varphi^t(T_{\mathbf{i}}) \delta_{\mathbf{i}}}{\sum_{\mathbf{i} \in A(k)} \varphi^t(T_{\mathbf{i}})}$$

where

$$A(k) = \{ \mathbf{i} : |\mathbf{i}| = k \text{ and } \exists \mathbf{j} \in \mathcal{A}^{\mathbb{N}} \text{ with } \mathbf{i}\mathbf{j} \in C \}$$

is the set of  $\mathbf{i}$  of length  $k$  that can be extended to be elements of  $C$  (as defined above). This way  $\mu_{n_k}$  is supported on the set of points that recur to the shrinking targets at times  $n_1, n_2, \dots, n_k$ . Define the probability measure  $\mu$  as the limit of a convergent subsequence of  $\{\mu_k\}$ .

Let  $|\mathbf{q}| = q$ . With condition (6) we have that

$$\begin{aligned}
 (10) \quad \mu_k[\mathbf{q}] &= \frac{\sum_{\substack{\mathbf{i} \in A(k) \\ \mathbf{i}|_q = \mathbf{q}}} \varphi^t(T_{\mathbf{i}})}{\sum_{\mathbf{i} \in A(k)} \varphi^t(T_{\mathbf{i}})} \leq \frac{\varphi^t(T_{\mathbf{q}}) \sum_{\substack{\mathbf{i} \in A(k) \\ \mathbf{i}|_q = \mathbf{q}}} \varphi^t(T_{\sigma^q(\mathbf{i})})}{D \sum_{\mathbf{i} \in A(k)} \varphi^t(T_{\mathbf{i}|_q}) \varphi^t(T_{\sigma^q(\mathbf{i})})} \\
 &= \frac{\varphi^t(T_{\mathbf{q}}) \sum_{\substack{\mathbf{i} \in A(k) \\ \mathbf{i}|_q = \mathbf{q}}} \varphi^t(T_{\sigma^q(\mathbf{i})})}{D \sum_{\mathbf{i} \in A(q)} \varphi^t(T_{\mathbf{i}}) \sum_{\substack{\mathbf{i} \in A(k) \\ \mathbf{i}|_q = \mathbf{q}}} \varphi^t(T_{\sigma^q(\mathbf{i})})} = \frac{\varphi^t(T_{\mathbf{q}})}{D \sum_{\mathbf{i} \in A(q)} \varphi^t(T_{\mathbf{i}})}
 \end{aligned}$$

for all  $k > q$ . We will now prove that the sequence  $\{n_k\}$  can be chosen so that there is some  $\kappa > 0$  such that

$$\Sigma_q := \sum_{\mathbf{i} \in A(q)} \varphi^t(T_{\mathbf{i}}) \geq \kappa$$

for all sufficiently large  $q$ . This will prove the lemma with  $H = (D\kappa)^{-1}$ .

Clearly,  $\Sigma_q$  decreases in intervals of the form  $n_k \leq q \leq n_k + \ell_{n_k}$ , so we first show that given a non-decreasing  $\iota : \mathbb{N} \rightarrow \mathbb{R}$ , we can choose  $\{n_k\}$  so that  $\Sigma_{n_k + \ell_{n_k}} \geq \iota(k)$  for all  $k$ .

Recall that  $P(s, \mathbf{j})$  is decreasing in  $s$ . Therefore, since  $t < s_0$ , there is some  $\varepsilon > 0$  such that

$$(11) \quad \sum_{|\mathbf{i}|=k} \varphi^t(T_{\mathbf{i}\mathbf{j}|_{\ell_k}}) \geq \exp(k\varepsilon)$$

for all  $k$  sufficiently large. Hence there is a number  $n_1$  such that

$$\Sigma_{n_1 + \ell_{n_1}} = \sum_{|\mathbf{i}|=n_1} \varphi^t(T_{\mathbf{i}\mathbf{j}|_{\ell_{n_1}}}) \geq \exp(n_1\varepsilon) \geq \iota(1).$$

Proceeding inductively, assume we have already chosen  $n_1, n_2, \dots, n_{k-1}$  such that  $\Sigma_{n_{k-1} + \ell_{n_{k-1}}} \geq \iota(k-1)$ . Then

$$\begin{aligned}
 \Sigma_{n_k + \ell_{n_k}} &= \sum_{|\mathbf{i}_1|=n_1} \sum_{|\mathbf{i}_2|=n_2 - n_1 - \ell_{n_1}} \cdots \sum_{|\mathbf{i}_k|=n_k - n_{k-1} - \ell_{n_{k-1}}} \varphi^t(T_{\mathbf{i}_1\mathbf{j}|_{\ell_{n_1}} \dots \mathbf{i}_k\mathbf{j}|_{\ell_{n_k}}}) \\
 &\stackrel{(6)}{\geq} \sum_{|\mathbf{i}_1|=n_1} \cdots \sum_{|\mathbf{i}_k|=n_k - n_{k-1} - \ell_{n_{k-1}}} D \varphi^t(T_{\mathbf{i}_1\mathbf{j}|_{\ell_{n_1}} \dots \mathbf{i}_{k-1}\mathbf{j}|_{\ell_{n_{k-1}}}}) \varphi^t(T_{\mathbf{i}_k\mathbf{j}|_{\ell_{n_k}}}) \\
 &= \Sigma_{n_{k-1} + \ell_{n_{k-1}}} D \sum_{|\mathbf{i}|=n_k - n_{k-1} - \ell_{n_{k-1}}} \varphi^t(T_{\mathbf{i}\mathbf{j}|_{\ell_{n_k}}}).
 \end{aligned}$$

Now, by the induction hypothesis,

$$\begin{aligned}
 &\geq \iota(k-1)D \left( \sum_{|\mathbf{i}|=n_{k-1}+\ell_{n_{k-1}}} \varphi^t(T_{\mathbf{i}}) \right)^{-1} \sum_{|\mathbf{j}|=n_k} \varphi^t(T_{\mathbf{i}\mathbf{j}|_{\ell_{n_k}}}) \\
 (12) \quad &\stackrel{(11)}{\geq} \left[ \iota(k-1)D \left( \sum_{|\mathbf{i}|=n_{k-1}+\ell_{n_{k-1}}} \varphi^t(T_{\mathbf{i}}) \right)^{-1} \right] \exp(n_k \varepsilon),
 \end{aligned}$$

as long as we choose  $n_k$  sufficiently large. Notice that the quantity in square brackets is a positive number only depending on our choices of  $n_1, \dots, n_{k-1}$ , so we can certainly choose  $n_k$  large enough that (12) exceeds  $\iota(k)$ , as claimed.

Now, suppose that we are in an interval of the form  $n_k + \ell_{n_k} \leq q \leq n_{k+1}$ , and  $q - n_k - \ell_{n_k} = \ell$ . Then (6) implies that

$$\Sigma_q \geq D \Sigma_{n_k + \ell_{n_k}} \sum_{\mathbf{i}=\ell} \varphi^t(T_{\mathbf{i}}).$$

Since  $t < s_0 < \dim E_{\mathbf{a}}$ , the expression  $\sum_{\mathbf{i}=\ell} \varphi^t(T_{\mathbf{i}})$  is bounded below by a constant that is uniform over  $\ell \in \mathbb{N}$ , which proves that there is some  $\kappa > 0$  such that  $\Sigma_q \geq \kappa$  for  $q$  sufficiently large, as wanted.

Returning to (10), the lemma is proved with  $H = (D\kappa)^{-1}$ .  $\square$

**Lemma 7.** *The image measure  $\pi_*\mu$  given by Lemma 6 has finite s-energy for almost all  $\mathbf{a}$ .*

*Proof.* Using [7, Lemma 3.1], [25, Proposition 3.1], and Fubini's theorem, the problem inside any ball of radius  $\rho$  reduces to the study of finiteness of

$$I = \iint_{R \times R} \frac{d\mu(\mathbf{i}) d\mu(\mathbf{k})}{\varphi^s(T_{\mathbf{i} \wedge \mathbf{k}})}.$$

Now, using (9), the fact that  $\mu$  is a probability measure, and  $t > s$

$$\begin{aligned}
 I &\leq \sum_{q=1}^{\infty} \sum_{|\mathbf{q}|=q} \sum_{i \neq k} \varphi^s(T_{\mathbf{q}})^{-1} \mu[\mathbf{q}i] \mu[\mathbf{q}k] \\
 &\leq Hm \sum_{q=1}^{\infty} \sum_{|\mathbf{q}|=q} \sum_{k=1}^m \frac{\varphi^t(T_{\mathbf{q}})}{\varphi^s(T_{\mathbf{q}})} \mu[\mathbf{q}k] \\
 &\leq Hm \sum_{q=1}^{\infty} \sigma_+^{q(t-s)} < \infty.
 \end{aligned}$$

Since  $\rho$  is arbitrary, the claim follow for almost all  $\mathbf{a}$ .  $\square$

*Proof of Theorem 1.* Fix a center  $\mathbf{j}$  and a sequence  $(\ell_k)$  of target sizes such that the limit  $\lim_{k \rightarrow \infty} \frac{1}{k} \log \varphi^s(T_{\mathbf{j}|\ell_k})$  exists. The almost-sure existence and uniqueness of the zero point  $s_0$  of the pressure follow from Lemma 3, and the estimate  $\dim_{\mathcal{H}} \tilde{R}(\pi(\mathbf{j})) \leq s_0$  from Lemma 4.

For the lower bound, let  $s < s_0$ . By Lemma 7 and the potential theoretic characterization of the Hausdorff dimension, since  $\mu$  is supported on  $C$ , we have  $\dim_{\mathcal{H}} \pi(C) \geq s$  almost surely. Approaching  $s_0$  along a sequence gives the lower bound for the dimension of  $\pi(C) \subset \tilde{R}(\pi(\mathbf{j}))$ .  $\square$

*Remark.* Notice that, the dimension of the recurring set is not affected by the set of points where  $\pi$  is not an injection, as long as this set has dimension smaller than  $s_0$ . This is the case for large classes of self-affine sets, for a Lebesgue-positive proportion of translation vectors  $\mathbf{a}$ , and in these cases there is the more geometric interpretation of §2.3 for the result of Theorem 1.

## 6. SOME FURTHER OBSERVATIONS

It is classical to study the shrinking target problem for an invariant measure. Unfortunately there is no canonical choice for a measure on a self-affine set. There is, however, a natural family of measures, called Gibbs measures, which we now define. In general, a Gibbs measure for the potential  $\varphi^t$ , or a **Gibbs measure at  $t$** , is a  $\sigma$ -invariant probability measure  $\mu$  on the symbolic space  $\mathcal{A}^{\mathbb{N}}$  satisfying

$$(13) \quad C^{-1} \leq \frac{\mu[\mathbf{q}]}{\varphi^t(T_{\mathbf{q}}) \exp(-|\mathbf{q}| \mathcal{P}(t))} \leq C.$$

Unfortunately, as demonstrated by [18, Example 6.4], Gibbs measures do not always exist. However, under the assumption (6) they do and, further, a Gibbs measure at the zero point of the pressure  $\mathcal{P}$  is a measure of maximal dimension, that is,

$$\dim_{\mathcal{H}}(\pi_*\mu) = \dim_{\mathcal{H}} E_{\mathbf{a}}$$

for almost all  $\mathbf{a} \in \mathbb{R}^{dm}$ . These facts were proved in [18, Theorem 5.2], [18, p. 8-9] and [17, Theorem 2.2]. Also see [15, Theorem 4.1]. This means that the Gibbs measure is a natural candidate as a measure to the size of the recurring set. Now we are in the position to formulate and prove the following corollary.

**Corollary 8** (Corollary to Theorem 2.1 of [5]). *Assume  $E$  is a self-affine set as in Theorem 1. Let  $(\ell_k) \subset \mathbb{N}$  be an increasing sequence tending to  $\infty$ , and  $\mathbf{j} \in \mathcal{A}^{\mathbb{N}}$ . Denote by  $\mu$  the Gibbs measure at the point*

$t$  where  $\mathcal{P}(t) = 0$ . Then the measure of the recurrence set  $\pi_*\mu(\tilde{R}(\pi(\mathbf{j})))$  is either 0 or 1 according to whether the sum

$$(14) \quad \sum_{k=1}^{\infty} \mu([\mathbf{j}_{\ell_k}])$$

converges or diverges.

*Proof.* First, the fact that the convergence of the sum (14) implies  $\pi_*\mu(\tilde{R}(\pi(\mathbf{j}))) = 0$  follows from the classical Borel–Cantelli lemma. This is the case since  $\pi_*\mu(\tilde{R}(\pi(\mathbf{j}))) = \mu(R(\mathbf{j}))$  and  $\mu$  is  $\sigma$ -invariant.

Now assume that the sum (14) diverges. Notice that while the definition of Gibbs measures in the symbolic space corresponding to self-affine sets does not necessarily coincide with the classical one, under the assumption (6) the measure  $\mu$  satisfies the Facts 1–3 in Section 3 of [5]. As these facts are all the knowledge on  $\mu$  needed to prove Theorem 2.1 of [5], it follows that  $\mu(R(\mathbf{j})) = \pi_*\mu(\tilde{R}(\pi(\mathbf{j})))$  has measure 1.  $\square$

We finish with two further questions related to Theorem 1.

- Will the claim of Theorem 1 remain true without condition (6)?
- What about recurrence under  $F$  to geometric sets  $B_k$ , for example to balls in  $\mathbb{R}^d$ ? What could be the counterpart of  $s_0$ ? One has to be a little careful when stating a result of this type since, as pointed out above, in the overlapping cases  $F$  is not even well-defined, and in most cases the corresponding symbolic problem becomes very hard to track. In our treatment it is possible that points fairly close to the target point  $\pi(\mathbf{j})$  on the fractal set are not considered to be recurring, if they are far away from  $\mathbf{j}$  in the symbolic space. Taking this kind of shortcut becomes impossible in the geometric case.

## REFERENCES

- [1] T. Bedford. Crinkly curves, Markov partitions and box dimensions in self-similar sets. *PhD Thesis, The University of Warwick*, 1984. [2](#)
- [2] Victor Beresnevich and Sanju Velani. A mass transference principle and the Duffin-Schaeffer conjecture for Hausdorff measures. *Ann. of Math. (2)*, 164(3):971–992, 2006. [3](#)
- [3] Rufus Bowen. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*. Lecture Notes in Mathematics, Vol. 470. Springer-Verlag, Berlin-New York, 1975. [3](#)
- [4] Yann Bugeaud, Stephen Harrap, Simon Kristensen, and Sanju Velani. On shrinking targets for  $\mathbb{Z}^m$  actions on tori. *Mathematika*, 56(2):193–202, 2010. [2](#)
- [5] N. Chernov and D. Kleinbock. Dynamical Borel-Cantelli lemmas for Gibbs measures. *Israel J. Math.*, 122:1–27, 2001. [1](#), [3](#), [13](#), [14](#)

- [6] K. J. Falconer. The dimension of self-affine fractals. II. *Math. Proc. Cambridge Philos. Soc.*, 111(1):169–179, 1992. [2](#)
- [7] Kenneth Falconer. The Hausdorff dimension of self-affine fractals. *Math. Proc. Cambridge Philos. Soc.*, 103(2):339–350, 1988. [2](#), [6](#), [12](#)
- [8] Bassam Fayad. Mixing in the absence of the shrinking target property. *Bull. London Math. Soc.*, 38(5):829–838, 2006. [2](#)
- [9] Andrew Ferguson, Thomas Jordan, and Michał Rams. Dimension of self-affine sets with holes. [3](#)
- [10] Richard Hill and Sanju L. Velani. The ergodic theory of shrinking targets. *Invent. Math.*, 119(1):175–198, 1995. [1](#)
- [11] Richard Hill and Sanju L. Velani. The shrinking target problem for matrix transformations of tori. *J. London Math. Soc. (2)*, 60(2):381–398, 1999. [1](#)
- [12] Richard Hill and Sanju L. Velani. A zero-infinity law for well-approximable points in Julia sets. *Ergodic Theory Dynam. Systems*, 22(6):1773–1782, 2002. [1](#)
- [13] John Hutchinson. Fractals and self-similarity. *Indiana Univ. Math. J.*, 30(5):713–747, 1981. [2](#)
- [14] Antti Käenmäki. On natural invariant measures on generalised iterated function systems. *Ann. Acad. Sci. Fenn. Math.*, 29(2):419–458, 2004. [10](#)
- [15] Antti Käenmäki and Henry W. J. Reeve. Multifractal analysis of Birkhoff averages for typical infinitely generated self-affine sets. *J. Fractal Geom.*, 1(1):83–152, 2014. [3](#), [13](#)
- [16] Antti Käenmäki and Pablo Shmerkin. Overlapping self-affine sets of Keakeya type. *Ergodic Theory Dynam. Systems*, 29(3):941–965, 2009. [6](#)
- [17] Antti Käenmäki and Markku Vilppolainen. Separation conditions on controlled Moran constructions. *Fund. Math.*, 200(1):69–100, 2008. [13](#)
- [18] Antti Käenmäki and Markku Vilppolainen. Dimension and measures on sub-self-affine sets. *Monatsh. Math.*, 161(3):271–293, 2010. [13](#)
- [19] J. Kurzweil. On the metric theory of inhomogeneous diophantine approximations. *Studia Math.*, 15:84–112, 1955. [2](#)
- [20] Bing Li, Bao-Wei Wang, Jun Wu, and Jian Xu. The shrinking target problem in the dynamical system of continued fractions. *Proc. Lond. Math. Soc. (3)*, 108(1):159–186, 2014. [2](#)
- [21] Curt McMullen. The Hausdorff dimension of general Sierpiński carpets. *Nagoya Math. J.*, 96:1–9, 1984. [2](#)
- [22] Tomas Persson and Michał Rams. On shrinking targets for piecewise expanding rational maps. available online at <http://arxiv.org/abs/1406.6785>. [2](#)
- [23] Tapio Rajala and M. Vilppolainen. Weakly controlled moran constructions and iterated function systems in metric spaces. to appear in *Illinois J. Math.* available online at <http://arxiv.org/abs/1003.0349>. [10](#)
- [24] Henry Reeve. Shrinking targets for countable Markov maps. available online at <http://arxiv.org/abs/1107.4736>. [2](#)
- [25] Boris Solomyak. Measure and dimension for some fractal families. *Math. Proc. Cambridge Philos. Soc.*, 124(3):531–546, 1998. [12](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YO10 5DD, UK

*E-mail address:* [henna.koivusalo@york.ac.uk](mailto:henna.koivusalo@york.ac.uk)



WESLEYAN UNIVERSITY, MIDDLETOWN, CT, USA

*E-mail address:* framirez@wesleyan.edu